

論文タイトル等	編者・著者	掲載情報 雑誌名, 掲載年; 巻(号): ページ番号
An association between serum uric acid levels/hyperuricemia and hypertension among 85,900 Japanese workers: Risk of hypertension.	Yokokawa H, Fukuda H, Suzuki A, Fujibayashi K, Naito T, Uehara Y, Nakayama A, Matsuo H, Sanada H, Jose PA, Miwa Y, Hisaoka T, Isonuma H.	J Clin Hypertens (Greenwich). 2015. 18:53-59.
Utility of 75-g oral glucose tolerance test results and hemoglobin a1c values for predicting the incidence of diabetes mellitus among middle-aged Japanese men -a large-scale retrospective cohort study performed at a single hospital	Fujibayashi K, Yokokawa H, Gunji T, Sasabe N, Okumura M, Iijima K, Haniu T, Hisaoka T, Fukuda H.	Intern Med. 2015;54(7):717-23.
Raltegravir and abacavir/lamivudine in Japanese treatment-naïve and treatment-experienced HIV-infected patients: a 48-week retrospective pilot analysis.	Suzuki A, Uehara Y, Saita M, Inui A, Isonuma H, Naito T	JJID 69: 33-38, 2015
Global distribution of polymorphisms associated with delayed Plasmodium falciparum parasite clearance following artemisinin treatment: genotyping of archive blood samples.	Murai K, Culleton R, Hisaoka T, Endo H, Mita T.	Parasitol Int 2015. 64: 267-73. PMID: 25449286
Relationship between causes of fever of unknown origin and inflammation markers: a multicenter collaborative retrospective study.	Naito T, Torikai K, Mizooka M, Mitsumoto F, Kanazawa K, Ohno S, Morita H, Ukimura A, Mishima N, Otsuka F, Ohyama Y, Nara N, Murakami K, Mashiba K, Akazawa K, Yamamoto K, Tanei M, Yamanouchi M, Senda S, Tazuma S, Hayashi J.	Intern Med, 2015; 54: 1989-1994.
Body mass index and waist circumference are independent risk factors for low vital capacity among Japanese participants of a health checkup: a single-institution cross-sectional study.	Goto Y, Yokokawa H, Fukuda H, Naito T, Hisaoka T, Isonuma H.	World Academy of Science, Engineering and Technology, 2015: 20: 108-115.
Seroprevalence, seroconversion, and toxoplasmosis risk factors among pregnant women in Taipei City, Taiwan.	Hung CS, Su HW, Lee YL, Weng HW, Wang YC, Naito T, Tsubouchi A, Wang GC, Fan CK	Jpn J Infect Dis, 2015; 68: 312-317.
Seroprevalence and Associated Risk Factors of Toxocariasis among College Students in Taipei City, Taiwan.	Fu CJ, Kao CY, Lee YL, Liao CW, Chen PC, Chuang TW, Wang YC, Chou CM, Huang YC, Naito T, Fan CK	Iran J Parasitol, 2015; 10: 482-189.
Age- and Sex-Specific Impact of Health Literacy on Healthy Lifestyle Characteristics among Japanese Residents in a Rural Community	Hirohide Yokokawa, Motoyuki Yuasa, Hironobu Sanada, Teruhiko Hisaoka, Hiroshi Fukuda	Health, 2015 ; 7(6) : DOI: 10.4236/health.2015.76081
Association Between Serum Uric Acid Levels/Hyperuricemia and Hypertension Among 85,286 Japanese Workers	Hirohide Yokokawa, Hiroshi Fukuda, Akihito Suzuki, Kazutoshi Fujibayashi, Toshio Naito, Yuki Uehara, Akiyoshi Nakayama, Hirotaka Matsuo, Hironobu Sanada, Pedro A. Jose, Yuichi Miwa, Teruhiko Hisaoka, Hiroshi Isonuma	The Journal of Clinical Hypertension, 2015; DOI: 10.1111/jch.12627
Achievement of Target Blood Pressure Levels among Japanese Workers with Hypertension and Healthy Lifestyle Characteristics Associated with Therapeutic Failure	Nagako Kudo, Hirohide Yokokawa, Hiroshi Fukuda, Hironobu Sanada, Yuichi Miwa, Teruhiko Hisaoka, Hiroshi Isonuma	PLOS ONE, 2015; DOI:10.1371/journal.pone.0133641
Edwardsiella tarda bacteremia. A rare but fatal water- and foodborne infection: Review of the literature and clinical cases from a single	Hirai Y, Asahata-Tago S, Ainoda Y, Fujita T, Kikuchi K.	Can J Infect Dis Med Microbiol. 2015 Nov-Dec;26(6):313-8. Review.
Candidemia Diagnosed from Peripheral Blood Smear: Case Report and Review of Literature 1954-2013.	Hirai Y, Asahata S, Ainoda Y, Fujita T, Miura H, Hizuka N, Kikuchi K.	Mycopathologia. 2015 Aug;180(1-2):111-6.
Fournier's gangrene caused by Listeria monocytogenes as the primary organism.	Asahata S, Hirai Y, Ainoda Y, Fujita T, Okada Y, Kikuchi K.	Can J Infect Dis Med Microbiol. 2015 Jan-Feb;26(1):44-6.
Risk factors for recurrent hospital-acquired Clostridium difficile infection in a Japanese university hospital.	Hikone M, Ainoda Y, Tago S, Fujita T, Hirai Y, Takeuchi K, Totsuka K.	Clin Exp Gastroenterol. 2015 Jul 14;8:191-6.
Gram-negative rod bacteremia after cardiovascular surgery: Clinical features and prognostic factors.	Tago S, Hirai Y, Ainoda Y, Fujita T, Kikuchi K.	J Microbiol Immunol Infect. 2015 Aug 4. pii: S1684-1182(15)00814-2.
Perianal tuberculosis: A case report and review of the literature.	Tago S, Hirai Y, Ainoda Y, Fujita T, Takamori M, Kikuchi K.	World J Clin Cases. 2015;16:3(9):848-52.
Decidual β -carotene-15,15'-oxygenase-1 and 2 (BCMO1,2) expression is increased in nitrofen model of congenital diaphragmatic hernia.	Takahashi H, Kutasy B, Pes L, Paradisi F, Puri P.	Pediatr Surg Int. 2015 Jan;31(1):37-43. doi: 10.1007/s00383-014-3621-8. Epub 2014 Oct
Kif7 expression is decreased in the diaphragmatic and pulmonary mesenchyme of nitrofen-induced congenital diaphragmatic hernia.	Takahashi T, Friedmacher F, Takahashi H, Hofmann AD, Puri P.	J Pediatr Surg. 2015 Jun;50(6):904-7. doi: 10.1016/j.jpedsurg.2015.03.058. Epub 2015 Apr 11.

論文タイトル等	編者・著者	掲載情報 雑誌名, 掲載年 ; 巻(号) : ページ番号
Seroprevalence of Toxoplasma gondii Infection among HIV-Infected Patients in Tokyo, Japan.	Hiromizu Takahashi, Akiko Tsubouchi, Kenji Murai, Masashi Yamanouchi, Mika Tanei, Rino Sakamoto, Yuki Uehara, Chia-Kwung Fan and Toshio Naito	Clin Res HIV/AIDS 1(2): 1007 (2014)
Prenatal administration of all-trans retinoic acid upregulates leptin signaling in hypoplastic rat lungs with experimental congenital diaphragmatic hernia.	Friedmacher F, Hofmann AD, Takahashi T, Takahashi H, Kutasy B, Puri P.	Pediatr Surg Int. 2014 Dec;30(12):1183-90. doi: 10.1007/s00383-014-3605-8. Epub 2014 Oct 21.
Disruption of copper-dependent signaling pathway in the nitrofen-induced congenital diaphragmatic hernia.	Takahashi T, Friedmacher F, Takahashi H, Hofmann AD, Puri P.	Pediatr Surg Int. 2015 Jan;31(1):31-5. doi: 10.1007/s00383-014-3620-9. Epub 2014 Oct 16.
Myogenin gene expression is not altered in the developing diaphragm of nitrofen-induced congenital diaphragmatic hernia.	Takahashi T, Friedmacher F, Takahashi H, Hofmann AD, Puri P.	Pediatr Surg Int. 2014 Sep;30(9):901-6. doi: 10.1007/s00383-014-3557-z. Epub
Evidence for decreased lipofibroblast expression in hypoplastic rat lungs with congenital diaphragmatic hernia.	Friedmacher F, Fujiwara N, Hofmann AD, Takahashi H, Gosemann JH, Puri P.	Pediatr Surg Int. 2014 Oct;30(10):1023-9. doi: 10.1007/s00383-014-3549-z. Epub
Prenatal retinoic acid increases lipofibroblast expression in hypoplastic rat lungs with experimental congenital diaphragmatic hernia.	Friedmacher F, Fujiwara N, Hofmann AD, Takahashi H, Alvarez LA, Gosemann JH, Puri P.	J Pediatr Surg. 2014 Jun;49(6):876-81; discussion 881. doi: 10.1016/j.jpedsurg.2014.01.017. Epub 2014 Feb 3.